Universal Statistics of Branched Flows

Jakob J. Metzger^{1,2}, Ragnar Fleischmann¹, Theo Geisel^{1,2}

¹Max-Planck-Institute for Dynamics and Self-Organization, Göttingen, Germany

²Department of Physics, University of Göttingen, Germany

Universal Statistics of Branched Flows

Jakob J. Metzger^{1,2}, Ragnar Fleischmann¹, Theo Geisel^{1,2}

¹Max-Planck-Institute for Dynamics and Self-Organization, Göttingen, Germany

²Department of Physics, University of Göttingen, Germany

FOR760

Cargèse, July 2010

Experimental observation of branched flow

Topinka *et al.*, Nature **410**, 183 (2001)

Motivation

- Branched flow is a universal phenomenon of 2D Hamiltonian flows in weak, correlated disorder potentials
- Many flows can approximately be described by Hamiltonian flows and suffer from small perturbations by a random potential

- Examples:

- Branched flow in semiconductor devices (with magnetic fields*)

Topinka *et al.*, Nature **410**, 183 (2001)

Aidala *et al.*, Nat. Phys. **3** (2007) *

Jura *et al.*, Nat. Phys. **3** (2007)

- Freak waves in microwave cavities

Höhmman *et al.*, PRL **104** (2010)

- Rogue sea waves

Heller *et al.*, J. Geophys. Res. **113** (2008)

Müller *et al.*, Oceanography, **18** (2005)

- Branching of sound waves

Wolfson, Tomsovic, J. Acous. Soc. Am. **109** (2001)

⋮

Talk Outline

- Motivation
- Intro: Branched Flow and Caustics
- Branch/caustics statistics
Approximations and Results
- Conclusion/Outlook

Caustics and branched flow

$$\langle V(\mathbf{r}')V(\mathbf{r}' + \mathbf{r}) \rangle = \epsilon^2 c(|\mathbf{r}|/\ell_c)$$

$$\mathbf{r} = (x, y)$$

$$H = \mathbf{p}^2/2 + V(\mathbf{r})$$

Caustics and branched flow

$$\langle V(\mathbf{r}')V(\mathbf{r}' + \mathbf{r}) \rangle = \epsilon^2 c(|\mathbf{r}|/\ell_c)$$

$$\mathbf{r} = (x, y)$$

$$H = \mathbf{p}^2/2 + V(\mathbf{r})$$

Caustics and branched flow

$$\langle V(\mathbf{r}')V(\mathbf{r}' + \mathbf{r}) \rangle = \epsilon^2 c(|\mathbf{r}|/\ell_c)$$

$$\mathbf{r} = (x, y)$$

$$H = \mathbf{p}^2/2 + V(\mathbf{r})$$

Caustics and branched flow

$$\langle V(\mathbf{r}')V(\mathbf{r}' + \mathbf{r}) \rangle = \epsilon^2 c(|\mathbf{r}|/\ell_c)$$

$$\mathbf{r} = (x, y)$$

$$H = \mathbf{p}^2/2 + V(\mathbf{r})$$

Caustics and branched flow

$$\langle V(\mathbf{r}')V(\mathbf{r}' + \mathbf{r}) \rangle = \epsilon^2 c(|\mathbf{r}|/\ell_c)$$

$$\mathbf{r} = (x, y)$$

$$H = \mathbf{p}^2/2 + V(\mathbf{r})$$

Caustics and branched flow

$$\langle V(\mathbf{r}')V(\mathbf{r}' + \mathbf{r}) \rangle = \epsilon^2 c(|\mathbf{r}|/\ell_c)$$

$$\mathbf{r} = (x, y)$$

$$H = \mathbf{p}^2/2 + V(\mathbf{r})$$

Caustics and branched flow

How many branches exist on average per unit length?

History

- Kravtsov, Sov. Phys. JETP **28** (1969)

- Berry & Upstill, Progr. in Optics **18** (1980)

Fig. 7.1. Clusters of cusps in the caustics show a corrugated wavefront.

- Kulkarny & White, Phys. Fluids, **25**, (1982)
- White & Fornberg, J. Fluid Mech. **335** (1998)

mean first time to caustic $\langle t_{caustic} \rangle \propto \epsilon^{-2/3} \ell_c$
 scaling of mean free path $\ell_{mfp} \propto \epsilon^{-2} \ell_c$

- Kaplan, Phys. Rev. Lett. **89** (2002)

Branch counting

How many branches transverse to flow exist on average as a function of distance?

$$N_b(t) = \frac{1}{2} \lim_{L \rightarrow \infty} \frac{1}{L} \left\langle \int_0^L dy_0 \delta(m(y_0, t)) |\partial_{y_0} m(y_0, t)| \right\rangle \quad [1]$$

$$= \frac{1}{2} \lim_{L \rightarrow \infty} \frac{1}{L} \left\langle \int_0^L dy_0 \frac{\delta(t - t_c)}{|\dot{m}(y_0, t)|} |\partial_{y_0} m(y_0, t)| \right\rangle$$

$P_c(t)$ probability to caustic along trajectory [2]

$m(t) \equiv \partial y(t)/\partial y(0)$ entry of the stability matrix $m_{ij}(t) \equiv \partial \gamma_i(t)/\partial \gamma_j(0)$

$n(t) \equiv \partial_{y_0} m(t)$ entry of the extended stability tensor $n_{ijk}(t) \equiv \partial^2 \gamma_i(t)/\partial \gamma_j(0) \partial \gamma_k(0)$

[1] Berry & Upstill, Progr. Opt. **18**, 257 (1980)

[2] White & Fornberg, J. Fluid Mech. **355**, 113–138 (1998)

Statistics of the stability matrix and tensor

- Coupled differential equations for entries of the stability matrix and extended stability tensor
- Nonlinear, multiplicative noise
- Obtained closed set of 11 linear differential equations for the moments

Approximations

Approximations

- short-term solution: $N_b(t) \approx c_1 (\sigma_2/\sigma_1) t_0 P_c(t) g(t/t_0) \quad t \ll t_0$

$$\sigma_1^2 = \frac{1}{2} \int_{-\infty}^{\infty} dx \left. \frac{\partial^4 c(x, y)}{\partial y^4} \right|_{y=0}$$

$$\sigma_2^2 = -\frac{1}{2} \int_{-\infty}^{\infty} dx \left. \frac{\partial^6 c(x, y)}{\partial y^6} \right|_{y=0}$$

$$t_0 \propto \sigma_1$$

Approximations

- short-term solution: $N_b(t) \approx c_1 (\sigma_2/\sigma_1) t_0 P_c(t) g(t/t_0) \quad t \ll t_0$
- long-term solution: $N_b(t) \approx c_2 (\sigma_2/\sigma_1) t_0 P_c(t) e^{\lambda(t/t_0)} \quad t \gg t_0$

$$\sigma_1^2 = \frac{1}{2} \int_{-\infty}^{\infty} dx \left. \frac{\partial^4 c(x, y)}{\partial y^4} \right|_{y=0}$$
$$\sigma_2^2 = -\frac{1}{2} \int_{-\infty}^{\infty} dx \left. \frac{\partial^6 c(x, y)}{\partial y^6} \right|_{y=0}$$
$$t_0 \propto \sigma_1$$

Approximations

- short-term solution: $N_b(t) \approx c_1 (\sigma_2/\sigma_1) t_0 \underbrace{P_c(t) g(t/t_0)}_{f_1(t/t_0)} \quad t \ll t_0$
- long-term solution: $N_b(t) \approx c_2 (\sigma_2/\sigma_1) t_0 \underbrace{P_c(t) e^{\lambda(t/t_0)}}_{f_2(t/t_0)} \quad t \gg t_0$

$$\sigma_1^2 = \frac{1}{2} \int_{-\infty}^{\infty} dx \left. \frac{\partial^4 c(x, y)}{\partial y^4} \right|_{y=0}$$

$$\sigma_2^2 = -\frac{1}{2} \int_{-\infty}^{\infty} dx \left. \frac{\partial^6 c(x, y)}{\partial y^6} \right|_{y=0}$$

$$t_0 \propto \sigma_1$$

Universal branch statistics

ϵ (%)	ℓ_c	$c(r)$		ϵ (%)	ℓ_c	$c(r)$		ϵ (%)	ℓ_c	$c(r)$	
+	4	0.1	I	▽	4	0.08	II	*	2	0.1	IV
*	8	0.1	I	○	6	0.08	II	▽	4	0.1	IV
◇	4	0.08	I	◇	4	0.06	II	◇	6	0.1	V
△	4	0.06	I	▷	2	0.01	II	◇	4	0.08	V
×	2	0.01	I	+	4	0.1	III	▽	2	0.1	VI
△	6	0.1	I (2D)	◇	2	0.08	III	◇	4	0.12	VI

- (I) $c(r) = \epsilon^2 \exp(-r^2/\ell_c^2)$
- (II) $c(r) = \epsilon^2 \operatorname{sech}(r/\ell_c)$
- (III) – (VI) ($\alpha = 1, 2, 3, 4$) $c(r) = \epsilon^2 (1 + r^2/\ell_c^2)^{-\alpha}$

Conclusions

- Branched flow is a widespread phenomenon of flows in weak, correlated random potentials on scales smaller than mean free path

- Branch statistics can now be calculated for all distances from source and for a wide range of correlation functions

- Universal scaling of branch statistics