Photons, dust, and honey bees

Diederik Wiersma

European Lab. For Non-linear Spectroscopy (LENS)
INO-CNR, Univ. of Florence

Photons, dust, and honey bees

Diederik Wiersma

European Lab. For Non-linear Spectroscopy (LENS)

INO-CNR, Univ. of Florence

www.complexphotonics.org

Micro and nano photonics group

European Lab. For Non-linear Spectroscopy (LENS)
INO-CNR, Univ. of Florence

www.complexphotonics.org

Pierre Barthelemy
Matteo Burresi
Francesca Intonti
Rajesh Kumar
Iacopo Malfanti
Vitantonio Matarazzo

Camilla Parmeggiani
Francesco Riboli
Romolo Savo
Radha Vivekananthan
Kevin Vynck
Jacopo Berlototti
Silvia Vignolini

Stefano Cavalieri
Marcello Colocci
Massimo Gurioli
Stefano Lepri
Roberto Livi
Roberto Righini

Optics of complex systems

Light transport in dielectric materials

Disordered

Powders
Porous materials

Partially ordered

Liquid crystals
Quasi crystals

Ordered

Photonic
crystals

Understanding light propagation

Understanding light propagation

Single scattering

Understanding light propagation

Photonic crystal

Understanding light propagation

Photonic glass

Disordered materials

Physics Today, August issue (2009)

Random materials

Light waves perform random walk

Average step ℓ (mean free path)

→ Diffusion process: $\langle x^2 \rangle = D \cdot t$ with $D = \frac{1}{3} v \ell$

Scattering

Interference effects

Speckle pattern

Analogies electrons - photons

Schrödinger's equation $[-\frac{\hbar^2}{2m}\nabla^2 + V(\vec{r})]\Psi(\vec{r}) = E\Psi(\vec{r})$

Maxwell's equations $[-\nabla^2 - \frac{\omega^2 \epsilon(\vec{r})}{c^2}]\vec{E}(\vec{r}) = \frac{\omega^2 \epsilon_0}{c^2}\vec{E}(\vec{r})$

Anderson localization

Very strongly scattering materials

$$k \ell < 1$$

$k \ell$: disorder parameter

Nature 390, 671 (1997)

Nature Physics 4, 359 (2008)

Localization in 1D

Random Stack of dielectric multilayers

Internal intensity distribution

Differences 1D vs 3D

In 1D: always localization for any disordered system

(light waves, sound, Schrödinger waves)

In 3D: phase transition in disorder parameter

k : wavevector, ℓ : mean free path

$k\ell \leq 1$
→

Lévy flights

Processes with Lévy statistics

Stock market fluctuations

Animal foraging

Human travel

Turbulent flow

Stable distributions

- Linear combination of elements remains in distribution
- Gaussian is limiting case with finite mean
- Rest has heavy tail asymptotic behaviour:

$$P(z) : \frac{1}{1 + z^{\alpha+1}}$$

$\alpha = 1$ Cauchy distribution

- Infinite variance and (for $\alpha \leq 1$) also infinite mean

Gaussian random walk

Δx from distribution with finite average ℓ and variance

Central limit theorem

$\sigma^2 \propto D \cdot t$ Gaussian distribution

Diffusion process: $\langle x^2 \rangle = D \cdot t$ with $D = \frac{1}{3} v \ell$

Lévy walk

Δx from stable distribution

Generalized central
limit theorem

Generalized
diffusion process: $\langle x^2 \rangle = D \cdot t^\gamma$ with: $\gamma = 3 - \alpha$ for $1 \leq \alpha < 2$

$\gamma = 1$: Normal diffusion $\gamma > 1$: Super diffusion

Lévy walks for light waves

Lévy walk for light

- Control step size $\langle z \rangle = \frac{1}{\sigma \cdot n}$ σ : cross section
 n : density
- Fractal particle size distribution?
→ does not work due to Rayleigh (Mie) scattering
- Engineer local particle density!

Lévy glass

TiO₂ nanoparticles + Glass Spheres + Sodium Silicate (liquid glass)

Glass and glass: index matched

Glass Spheres: introduce the density fluctuations

Diameter distribution:

$$P(d) \sim \frac{1}{d^{2+\alpha}}$$

(d between 5 and 650 μm)

Lévy glass

Lévy flight from multiple scattering on TiO₂ particles

Nature 453, 498 (May 22, 2008)

Sample design

Diameter distribution voids: $P(d) \sim \frac{1}{d^{2+\alpha}}$

Step length distribution: $P(z) \sim \frac{1}{1+z^{\alpha+1}}$

Super diffusion: $\langle x^2 \rangle = D \cdot t^\gamma$ with: $\gamma = 3 - \alpha$ for $1 \leq \alpha < 2$

Monte Carlo simulations

Monte Carlo simulations

Free space Lévy flight with step length:

$$P(z) \sim \frac{1}{1 + z^{\alpha+1}}$$

Monte Carlo simulations

Fixed geometry (quenched disorder)

Placing spheres in a space

Placing spheres in a space

Monte Carlo simulations

Random walk with quenched disorder

Quenching effects

Quenched vs annealed disorder

Experimental observations

Lévy glass

TiO₂ nanoparticles + Glass Spheres + Sodium Silicate (liquid glass)

TiO₂ nanoparticles + Glass Spheres + Polymer matrix (n matched)

Super diffusion

Enhanced diffusive transmission

Generalised Ohm's law:

$$T = \frac{1}{1 + aL^{\alpha/2}}$$

Transmission profiles

Diffusive case: small fluctuations

Levy Case: enhanced fluctuations

Enhanced spreading

Diffusive Case: Almost-gaussian profile

Lévy Case: Cusped spatial profile

Summary

- Transport of light in random systems
- How to realize non-gaussian optical disorder: Levy glass
- Superdiffusion of light, possibility to study optical Levy flights
- Open questions:
 - Unknown properties of Lévy flights (e.g. finite-size effects)
 - Weak localization (coherent backscattering), strong localization, speckle correlations, etc..

Micro and nano photonics group

European Lab. For Non-linear Spectroscopy (LENS)
INO-CNR, Univ. of Florence

www.complexphotonics.org

Pierre Barthelemy
Matteo Burrese
Francesca Intonti
Rajesh Kumar
Iacopo Malfanti
Vitantonio Matarazzo

Camilla Parmeggiani
Francesco Riboli
Romolo Savo
Radha Vivekananthan
Kevin Vynck
Jacopo Berlototti
Silvia Vignolini

Stefano Cavalieri
Marcello Colocci
Massimo Gurioli
Stefano Lepri
Roberto Livi
Roberto Righini