Semiclassical aspects of (chaotic) quantum systems

Part 1: Fundamental ideas, S -matrix-based observables

Klaus Richter

University of Regensburg, Germany

Semiclassical aspects of

Semiclassical aspects of

(chaotic) quantum systems

Semiclassics 1: fundamental ideas,

S -matrix-based observables

(Klaus Richter, University of Regensburg)

Semiclassical aspects of

(chaotic) quantum systems

Semiclassics 2: theory of spectral statistics (Sebastian Müller)

Universality in chaotic quantum systems

Random Matrix Theory (RMT)

classical

quantum

Advanced Semiclassical Theory

Universality in chaotic quantum systems

Random Matrix Theory (RMT)

classical

quantum

Advanced Semiclassical Theory

key: **classical orbit correlations**

Correlations in classical dynamics

Integrable systems

- ⇒ dynamics organized on tori in phase space
- ⇒ families of orbits with the same length and classical action $S = \oint \mathbf{p} d\mathbf{q}$

Chaotic systems

Example: Chaotic billiard:

Poincaré surface of section:

⇒ ergodic dynamics

Chaotic systems

H. Poincaré (1892):

- periodic orbits are densely distributed ; $\#po \sim e^{\mu T}$
- periodic orbits provide clue to overall behavior

Chaotic systems

- ⇒ subtle correlations between periodic orbits
- ⇒ periodic orbits appear in multiplets !

periodic orbit doublets

Example: correlated periodic orbit pair in the hyperbola billiard:

M. Sieber

⇒ action difference $S_1 - S_2 \sim \epsilon^2 p^2 \rightarrow 0$

M. Sieber and KR (2001)

periodic orbit doublets

periodic orbit structures

po pairs

po bundles

Müller et al. (2009)

periodic orbit structures

periodic orbit webs

- ⇒ role in classical mechanics ?
- ⇒ semiclassical backbone of chaotic quantum systems !

Correlations in spectral statistics

Quantum dynamics at mesoscopic scales

wavepacket evolution in a chaotic confinement:

(A. Goussev)

- complex wavefunction morphology
- **semiclassical regime**: $\lambda_{\text{dB}} \ll \text{system size } \mathcal{L}$

Universality in quantum spectra

Typical level sequences (Bohigas et al. (1983))

Universality in quantum spectra

Distribution of level spacings following from **Random Matrix Theory (RMT)**

Odlyzko (1989)

⇒ RMT predicts universality of level fluctuations:
for interacting systems, Riemann zeros, **classically chaotic systems**

Bohigas, Giannoni, Schmit (1983)

Universality in chaotic quantum systems

Random Matrix Theory (RMT)

classical

quantum

Advanced Semiclassical Theory

classical orbit correlations $\Rightarrow$ RMT predictions
 $\Rightarrow$ go beyond RMT !

Spectral form factor

- Quantum density of states: $d(E) = \sum_n \delta(E - E_n) = \bar{d}(E) + d_{\text{osc}}(E)$

Spectral form factor

- Quantum density of states: $d(E) = \sum_n \delta(E - E_n) = \bar{d}(E) + d_{\text{osc}}(E)$
- Spectral form factor:

$$K(t) \sim \int_{-\infty}^{\infty} d\eta \left\langle d_{\text{osc}} \left(E + \frac{\eta}{2} \right) d_{\text{osc}} \left(E - \frac{\eta}{2} \right) \right\rangle_E \exp(2\pi i \eta t)$$

Spectral form factor

- Quantum density of states: $d(E) = \sum_n \delta(E - E_n) = \bar{d}(E) + d_{\text{osc}}(E)$
- Spectral form factor:

$$K(t) \sim \int_{-\infty}^{\infty} d\eta \left\langle d_{\text{osc}} \left(E + \frac{\eta}{2} \right) d_{\text{osc}} \left(E - \frac{\eta}{2} \right) \right\rangle_E \exp(2\pi i \eta t)$$

$$\tau = T/T_H \quad ; \quad T_H = 2\pi\hbar \bar{d}(E)$$

chaotic system with time reversal symmetry
(quartic oscillator)

RMT form factor of the Gaussian Orthogonal Ensemble (GOE)

$$K^{\text{GOE}}(\tau) = 2\tau - \tau \log(1 + 2\tau) = 2\tau - 2\tau^2 + \dots \quad \text{for } \tau < 1$$

Form factor: Semiclassical evaluation

- Gutzwiller trace formula: coherent sum over periodic orbits

$$d_{\text{osc}}(E) = \frac{1}{\pi\hbar} \text{Re} \sum_{\gamma} A_{\gamma} e^{iS_{\gamma}(E)/\hbar} \quad ; \quad S_{\gamma}(E) \gg \hbar$$

Form factor: Semiclassical evaluation

- Gutzwiller trace formula: coherent sum over periodic orbits

$$d_{\text{osc}}(E) = \frac{1}{\pi\hbar} \text{Re} \sum_{\gamma} A_{\gamma} e^{iS_{\gamma}(E)/\hbar} \quad ; \quad S_{\gamma}(E) \gg \hbar$$

- Semiclassical representation of the form factor:

$$K(\tau) \sim \sum_{\gamma, \gamma'} \left\langle A_{\gamma} A_{\gamma'}^* e^{i(S_{\gamma} - S_{\gamma'})/\hbar} \delta \left(T - \frac{T_{\gamma} + T_{\gamma'}}{2} \right) \right\rangle_E$$

- non-vanishing contributions from pairs of correlated orbits
- diagonal approximation**: $\gamma = \gamma'$ (Berry (1985))

$$K(\tau) \simeq \frac{2}{T_{\text{H}}} \sum_{\gamma} |A_{\gamma}|^2 \delta(T - T_{\gamma})$$

- Sum rule (Hannay, Ozorio de Almeida (1985))

$$\sum_{\gamma} |A_{\gamma}|^2 \delta(T - T_{\gamma}) \simeq T \quad \Rightarrow$$

$$K(\tau) \approx 2\tau$$

Form factor: Off-diagonal contributions

- leading-order loop contribution:
- action difference:

$$\Delta S(\varepsilon) \simeq p^2 \varepsilon^2 / (2m\lambda)$$

- contribution to form factor:

Sieber and KR (2001)

$$K_{\text{off}}(\tau) \sim \text{Re} \int_0^\pi d\varepsilon \sum_{\gamma} |A_{\gamma}|^2 e^{i\Delta S(\varepsilon)/\hbar} \delta(T - T_{\gamma}) P(\varepsilon, T)$$

- density of crossings $P(\varepsilon, T)$?

Crossing distribution

$$P(\varepsilon, T) ?$$

Crossing distribution

$$T = NT_0 \quad ; \quad P(\varepsilon, T) \sim N^2 \sim T^2$$

Crossing distribution

$$P(\varepsilon, T) \sim \sin \varepsilon T^2$$

Crossing distribution

but: minimal loop time: $c = \epsilon e^{\lambda T_{\text{enc}}(\epsilon)/2} \quad ; \quad c \sim \pi$

Crossing distribution

$$P(\varepsilon, T) \sim \sin \varepsilon \left[T - 2T_{\text{enc}}(\varepsilon) \right]^2, \quad T_{\text{enc}} = \frac{2}{\lambda} \ln \frac{c}{\varepsilon}$$

Form factor: Off-diagonal contributions

- leading-order loop contribution:
- action difference:

$$\Delta S(\varepsilon) \simeq p^2 \varepsilon^2 / (2m\lambda)$$

- contribution to form factor:

Sieber, KR (2001)

$$K_{\text{off}}(\tau) \sim \text{Re} \int_0^\pi d\varepsilon \sum_{\gamma} |A_{\gamma}|^2 e^{i\Delta S(\varepsilon)/\hbar} \delta(T - T_{\gamma}) P(\varepsilon, T)$$

- density of crossings $P(\varepsilon, T)$?

$$P(\varepsilon, T) = \frac{v^2}{2\pi A} \sin \varepsilon [T - 2T_{\text{enc}}(\varepsilon)]^2$$

$$\Rightarrow K_{\text{off}}(\tau) \approx -2\tau^2$$

form factor: semiclassical evaluation

$$\begin{aligned} K^{\text{GOE}}(\tau) &= 2\tau - \tau \log(1 + 2\tau) \\ &= 2\tau - 2\tau^2 + \dots \end{aligned} \quad \text{for } \tau = T/T_H < 1$$

- phase space approaches:
D. Spehner (2003); M. Turek and KR (2003);
S Heusler, S Müller, P Braun, F Haake (2004)
- periodic orbit theory for the
complete RMT form factor
Müller, Heusler, Altland, Braun, Haake (2007)

form factor: semiclassical evaluation

$$\begin{aligned}
 K^{\text{GOE}}(\tau) &= 2\tau - \tau \log(1 + 2\tau) \\
 &= 2\tau - 2\tau^2 + \dots
 \end{aligned}
 \quad \text{for } \tau = T/T_H < 1$$

- phase space approaches:
D. Spehner (2003); M. Turek and KR (2003);
S Heusler, S Müller, P Braun, F Haake (2004)
- periodic orbit theory for the complete RMT form factor
Müller, Heusler, Altland, Braun, Haake (2007)

form factor: semiclassical evaluation

$$\begin{aligned} K^{\text{GOE}}(\tau) &= 2\tau - \tau \log(1 + 2\tau) \\ &= 2\tau - 2\tau^2 + \dots \end{aligned} \quad \text{for } \tau = T/T_H < 1$$

- phase space approaches:
D. Spehner (2003); M. Turek and KR (2003);
S Heusler, S Müller, P Braun, F Haake (2004)
- periodic orbit theory for spectral statistics
→ Sebastian Müller's lecture !

form factor for the diffusive case

- quantum (impurity) scattering takes role of encounter dynamics
- diagonal contributions sufficient
- relation between $K(t)$ and return probability $p(t)$:

$$K(t) = \frac{2}{(2\pi\hbar)^2} \frac{d\Omega}{dE} t p(t)$$

- diffusive case: $p(t) \sim (4\pi Dt)^{-N/2}$

$$K^{\text{diff}}(t) = \frac{2\Omega_x}{(2\pi\hbar)^2 (4\pi D)^{N/2}} t^{1-N/2}$$

Argaman, Imry, Smilansky (1993)

form factor for the diffusive case

- quantum (impurity) scattering takes role of encounter dynamics
- diagonal contributions sufficient
- relation between $K(t)$ and return probability $p(t)$:

$$K(t) = \frac{2}{(2\pi\hbar)^2} \frac{d\Omega}{dE} t p(t)$$

- diffusive case: $p(t) \sim (4\pi Dt)^{-N/2}$

$$K^{\text{diff}}(t) = \frac{2\Omega_x}{(2\pi\hbar)^2 (4\pi D)^{N/2}} t^{1-N/2}$$

Argaman, Imry, Smilansky (1993)

Semiclassical S -matrix

S-matrix based observables

- Wigner time delay:

$$\tau_W(E) = -\frac{i\hbar}{N} \text{Tr} \left[S^\dagger(E) \frac{dS(E)}{dE} \right]$$

S-matrix based observables

- Wigner time delay:

$$\tau_W(E) = -\frac{i\hbar}{N} \text{Tr} \left[S^\dagger(E) \frac{dS(E)}{dE} \right]$$

- mesoscopic quantum transport:

Landauer: "conductance is transmission": $G = (2e^2/h) \text{Tr} [tt^\dagger]$

- moments of transmission:

$$M_n = \text{Tr} \langle [tt^\dagger]^n \rangle$$

→ shot noise $\sim M_1 - M_2$

→ full counting statistics

- Andreev billiards

Semiclassical approach

- Scattering matrix elements as sum over classical paths:

$$S_{ba}(E) \sim \sum_{\gamma(a \rightarrow b)} A_{\gamma} e^{(i/\hbar) S_{\gamma}(E)}$$

Miller (1972) ; Blümel and Smilansky (1988)

Semiclassical approach

- Scattering matrix elements as sum over classical paths:

$$S_{ba}(E) \sim \sum_{\gamma(a \rightarrow b)} A_{\gamma} e^{(i/\hbar) S_{\gamma}(E)}$$

Miller (1972) ; Blümel and Smilansky (1988)

- Semiclassical treatment of correlation functions:

$$C(\epsilon, n) = \text{Tr} \langle [S^{\dagger}(E - \epsilon) S(E + \epsilon)]^n \rangle$$

trajectory sets with encounters contributing e.g. to $n = 3$:

treatment to leading order in $1/N$:
Berkolaiko, Harrison, Novaes, JPA (2008)

Ehrenfest time dependence:
Waltner, Kuipers, Richter, arXiv:1007:1595

Andreev billiards: semiclassics beyond RMT

Andreev reflection

A. Philip, Groningen

retro-reflection at
normal-superconductor-interface

Andreev Billiards

Andreev (retro-) reflection
at superconductor-billiard
interface
 $\Rightarrow$ periodic orbits

- 'Bohr-Sommerfeld' expression for DoS in billiard: $S_\gamma = (n + 1/2)h$

$$\Rightarrow d(E) \sim \int_0^\infty dL P(L) L \sum_n \delta \left(\frac{EL}{\hbar v_F} - \left(n + \frac{1}{2}\right)\pi \right)$$

Melsen, Brouwer, Frahm, Beenakker (1996) ; Lodder & Nazarov (1998)

- probe $P(L)$, i.e classical dynamics, through proximity effect !

Proximity effect on density of states

$$d(E) \sim \int_0^\infty dL P(L) L \sum_n \delta \left(\frac{EL}{\hbar v_F} - \left(n + \frac{1}{2}\right) \pi \right)$$

chaotic case: $P(L) \sim e^{-L/L_D}$

square billiard (rough walls)

gap in DoS

regular case: $P(L) \sim L^{-\alpha}$

square billiard

linear suppression of DoS

Ihra, Leadbeater, Vega & KR (2001)

Excitation gap in chaotic case

Random Matrix Theory:

hard gap

at Thouless energy: $E_T = \frac{\hbar}{2\tau_D}$

$$\epsilon = E/E_T$$

Melsen, Brouwer, Frahm
& Beenakker (1996)

'Bohr-Sommerfeld':

no gap

($\sim$ diagonal approximation)

Schomerus & Beenakker (1999);
Ihra, Leadbeater, Vega, KR & (2001)

Excitation gap in chaotic case

Random Matrix Theory:

hard gap

at Thouless energy: $E_T = \frac{\hbar}{2\tau_D}$

$$\epsilon = E/E_T$$

Melsen, Brouwer, Frahm
& Beenakker (1996)

'Bohr-Sommerfeld':

no gap

($\sim$ diagonal approximation)

Schomerus & Beenakker (1999);
Ihra, Leadbeater, Vega, KR & (2001)

way out: Ehrenfest time dependence

Lodder & Nazarov (1998)

Ehrenfest time effects

- Ehrenfest time τ_E : Chirikov et al. (1981)

separates wavepacket evolution following classical dynamics from time scales dominated by (random) wave interference

- $\mathcal{L} = \lambda_{\text{dB}} \exp(\lambda \tau_E)$; $\tau_E \sim |\ln \hbar|$
- short time correction to quantum universality

Excitation gap in chaotic case

Random Matrix Theory:

hard gap

at Thouless energy: $E_T = \frac{\hbar}{2\tau_D}$

$$\epsilon = E/E_T$$

universal regime

$$\tau = \tau_E/\tau_D \rightarrow 0$$

'Bohr-Sommerfeld':

no gap

($\sim$ diagonal approximation)

"classical" limit

$$\tau = \tau_E/\tau_D \rightarrow \infty$$

Excitation gap in chaotic case

Random Matrix Theory:

hard gap

at Thouless energy: $E_T = \frac{\hbar}{2\tau_D}$

$$\epsilon = E/E_T$$

universal regime

$$\tau = \tau_E/\tau_D \rightarrow 0$$

'Bohr-Sommerfeld':

no gap

($\sim$ diagonal approximation)

"classical" limit

$$\tau = \tau_E/\tau_D \rightarrow \infty$$

- semiclassical mechanism for hard RMT gap ?
- crossover: universal $\longrightarrow$ "classical" limit ?

Scattering approach to DoS

Scattering matrix:

$$\begin{aligned} \mathbf{S} &= \mathbf{S}_N \mathbf{S}_A \\ &= \begin{pmatrix} S(E) & 0 \\ 0 & S^*(-E) \end{pmatrix} \begin{pmatrix} 0 & -i \\ -i & 0 \end{pmatrix} \end{aligned}$$

↑ ↑ ↑
(e) part (h) part Andreev reflect.

Scattering approach to DoS

Scattering matrix:

$$\begin{aligned} \mathbf{S} &= \mathbf{S}_N \mathbf{S}_A \\ &= \begin{pmatrix} S(E) & 0 \\ 0 & S^*(-E) \end{pmatrix} \begin{pmatrix} 0 & -i \\ -i & 0 \end{pmatrix} \end{aligned}$$

$\uparrow$ $\uparrow$ $\uparrow$
 (e) part (h) part Andreev reflect.

Density of states in terms of scattering matrix:

$$d(\epsilon) \sim 1 + \frac{2}{M} \sum_{n=1}^{\infty} \frac{(-1)^n}{n} \frac{\partial}{\partial \epsilon} \text{Im} \left\{ \underbrace{\text{Tr} \left[S^\dagger \left(E_F - \frac{\epsilon \hbar}{2\tau_D} \right) S \left(E_F + \frac{\epsilon \hbar}{2\tau_D} \right) \right]^n}_{\equiv C(\epsilon, n)} \right\}$$

$$\epsilon = E/E_T$$

correlation function of n scattering matrices

Semiclassical approach

- Scattering matrix elements as sum over classical paths:

$$S_{ba}(E) \sim \sum_{\gamma(a \rightarrow b)} A_{\gamma} e^{(i/\hbar) S_{\gamma}(E)}$$

Semiclassical approach

- Scattering matrix elements as sum over classical paths:

$$S_{ba}(E) \sim \sum_{\gamma(a \rightarrow b)} A_{\gamma} e^{(i/\hbar) S_{\gamma}(E)}$$

- Correlation functions $C(\epsilon, n) = \text{Tr}[S^{\dagger}(-\epsilon)S(\epsilon)]^n$
in terms of sums over $2n$ paths
- trajectory structures of interfering paths ($n = 3$):

Evaluation of diagrams

- apply diagrammatic rules: (Heusler et al. (2006))
 - extend work on full counting statistics (Berkolaiko, Harrison & Novaes (2008))
- **generating function of correlators** (which includes all diagrams):

$$G(\epsilon, r) = \sum_{n=1}^{\infty} r^{n-1} C(\epsilon, n)$$

Evaluation of diagrams

- DoS in terms of the generating function:

$$\begin{aligned} H(\epsilon, r) &= \frac{1}{r} \frac{\partial}{\partial \epsilon} \int G(\epsilon, r) dr = \sum_{n=1}^{\infty} \frac{r^{n-1}}{n} \frac{\partial C(\epsilon, n)}{\partial \epsilon} \\ d(\epsilon) &= 1 - 2\text{Im } H(\epsilon, -1) \end{aligned}$$

- DoS given by solutions of a cubic equation

Evaluation of diagrams

- DoS in terms of the generating function:

$$H(\epsilon, r) = \frac{1}{r} \frac{\partial}{\partial \epsilon} \int G(\epsilon, r) dr = \sum_{n=1}^{\infty} \frac{r^{n-1}}{n} \frac{\partial C(\epsilon, n)}{\partial \epsilon}$$
$$d(\epsilon) = 1 - 2\text{Im } H(\epsilon, -1)$$

- DoS given by solutions of a cubic equation

This yields the RMT result

Kuipers, Waltner, Petitjean, Berkolaiko, KR,
Phys. Rev. Lett. (2010)

Finite Ehrenfest time

RMT limit:

$$\tau = \tau_E / \tau_D \rightarrow 0$$

hard gap

"classical" limit:

$$\tau = \tau_E / \tau_D \rightarrow \infty$$

no gap

crossover: universal RMT $\rightarrow$ "classical" limit ?

Ehrenfest time dependence of the gap width?

Finite Ehrenfest time

each encounter lasts an Ehrenfest time τ_E :

increase of τ_E :
forming diagrams with
encounters is less likely

increase of τ_E :
possibility increases for diagrams
with correlated trajectories along
their whole length

$$C(\epsilon, \tau, n) = C(\epsilon, n) e^{-\tau(1-in\epsilon)} + \frac{1 - e^{-\tau(1-in\epsilon)}}{1 - in\epsilon} \quad \tau = \tau_E / \tau_D$$

see also: Whitney & Jacquod (2006); Brouwer & Rahav (2006) for $n \leq 3$

Finite Ehrenfest time

- intermediate Ehrenfest times: $\tau = \tau_E / \tau_D = 2$

Finite Ehrenfest time

- intermediate Ehrenfest times: $\tau = \tau_E / \tau_D = 2$

- second gap at 'Ehrenfest energy' $E_E = \pi \hbar / 2 \tau_E$!

Kuipers, Waltner, Petitjean, Berkolaiko, KR, Phys. Rev. Lett. (2010)

Kuipers, Engl, Berkolaiko, Petitjean, Waltner, KR, arXiv:1004.1327

Summary and outlook

Advanced semiclassics can:

- explain universality of chaotic quantum systems
- go beyond random matrix theory

Summary and outlook

Advanced semiclassics can:

- explain universality of chaotic quantum systems
- go beyond random matrix theory
- tackle observables in the mesoscopic regime:
 - ▶ superconducting-normal conducting hybrids
 - ▶ quantum decay, time delay
 - ▶ fidelity
 - ▶ quantum transport (weak and strong localization)
- treat multi-component wave (Green) function structures:
 - ▶ spin-orbit interaction in low-dimensional semiconductors
 - ▶ pseudospin- and valley-degrees of freedom in graphene
 - ▶ Luttinger hamiltonian for holes

Thanks to:

- Jack Kuipers, Cyril Petitjean, Daniel Waltner, Thomas Engl, Gregory Berkolaiko
- Deutsche Forschungsgemeinschaft (FOR 760, GRK 638), Alexander von Humboldt Foundation