Light emission, local density of states and C_0 speckle correlation in strongly scattering media

Romain Pierrat, Alexandre Cazé and Rémi Carminati

Institut Langevin, ESPCI ParisTech, Paris

Light emission, local density of states and C_0 speckle correlation in strongly scattering media

Romain Pierrat, Alexandre Cazé and Rémi Carminati

Institut Langevin, ESPCI ParisTech, Paris

Summer school "Mesoscopic Physics in Complex Media"

1) Introduction

Computation of the **imaginary part of the Green function** at the position of the source in a random medium related to

- the **spontaneous decay rate** of the emitter;
- the **Local Density Of States (LDOS)**;
- the C_0 **intensity correlation**.

A study related to fundamental questions in mesoscopic physics as the ability

- to **image** local optical properties deep inside in the system;
- to **control** light emission by multiple scattering.

2) Physical and numerical model [1, 2]

■ Scattersers: N dipoles of polarisability

$$\alpha(\omega) = \frac{-3\pi c^3 \gamma}{\omega^3 (\omega - \omega_0 + i\gamma/2)}$$

■ Emitters: 1 source of decay rate ($\propto$ LDOS)

$$\frac{\Gamma}{\Gamma_0} = \frac{\rho}{\rho_0} = 1 + \frac{6\pi}{k} \Im \left[\text{Tr} \vec{\mathbf{S}}(\mathbf{r}_0, \mathbf{r}_0, \omega) \right]$$

$$\text{with } \vec{\mathbf{S}}(\mathbf{r}_0, \mathbf{r}_0, \omega) = \alpha(\omega) k^2 \sum_{k=1}^N \vec{\mathbf{G}}_0(\mathbf{r}_0, \mathbf{r}_k) \vec{\mathbf{G}}_k,$$

$$\text{and } \vec{\mathbf{G}}_j = \vec{\mathbf{G}}_0(\mathbf{r}_j, \mathbf{r}_0) + \alpha(\omega) k^2 \sum_{k=1, k \neq j}^N \vec{\mathbf{G}}_0(\mathbf{r}_j, \mathbf{r}_k) \vec{\mathbf{G}}_k.$$

4) LDOS and C_0 speckle correlation [3]

■ LDOS fluctuations and C_0 speckle correlation are equal [4]:

C_0 is of infinite range and a non-universal intensity correlation function [5]:

$$C_0 = \frac{\langle \rho^2(\mathbf{r}_0, \omega) \rangle}{\langle \rho(\mathbf{r}_0, \omega) \rangle^2} - 1 = \frac{\text{Var}[\rho(\mathbf{r}_0, \omega)]}{\langle \rho(\mathbf{r}_0, \omega) \rangle^2}$$

Expression valid for **all transport regime** for a non-absorbing medium.

■ LDOS distribution $P(\Gamma)$:

- ① $\Gamma < \Gamma_0$: collective interaction (multiple scattering)
- ② $P(\Gamma) \propto \Gamma^{-3/2}$: Near-field interaction with one scatterer
- ③ Near-field interaction with more than one scatterer sensitive to the correlation parameter f

$$N = 100, \omega = \omega_0 + \gamma, \omega_0 = 3 \times 10^{15} \text{ Hz}, \gamma = 1 \times 10^9 \text{ Hz}, R = 1.2 \mu\text{m}, R_0 = 0.05 \mu\text{m}, k\ell_B = 19, f = N(d_0/2)^3 / (R^3 - R_0^3)$$

■ Influence of regime ③ on C_0 :

- (a) Full calculation
  - (b) Two nearest neighbours taken into account
  - (c) Only the nearest neighbour taken into account
- C_0 is **strongly influenced** by the correlations of the disorder.

3) LDOS and multiple scattering [6]

■ LDOS distribution $P(\Gamma)$ and probability $P(\Gamma < \Gamma_0)$:

Possibility of $\Gamma < \Gamma_0$ due to collective interactions (interferences)

$P(\Gamma < \Gamma_0)$ can reach 50% depending on parameters

$$N = 500, \omega = \omega_0 \Rightarrow \lambda = 1.88 \mu\text{m}, R_0 = 0.36 \mu\text{m}, d_0 = 0.12 \mu\text{m}, f \leq 2\%$$

■ Average LDOS $\langle \Gamma \rangle$:

- No correlation in position
- No recurrent scattering
- Data described with

$$\epsilon_{LL} = 1 + \frac{\rho\alpha}{1 - \rho\alpha/3}$$

- Deviation from the independent scattering approximation

- No correlation in position
- **Signature of recurrent scattering**
- Data fitted numerically with

$$\epsilon_{eff} \neq \epsilon_{LL}$$

References

- [1] L. S. Froufe-Pérez, R. Carminati, and J. J. Sáenz. Fluorescence decay rate statistics of a single molecule in a disordered cluster of nanoparticles. *Phys. Rev. A*, 76:013835, 2007.
- [2] L. S. Froufe-Pérez and R. Carminati. Lifetime fluctuations of a single emitter in a disordered nanoscopic system: The influence of the transition dipole orientation. *Phys. Stat. Sol. (a)*, 205(6):1258, 2008.
- [3] A. Cazé, R. Pierrat, and R. Carminati. Near-field interactions and nonuniversality in speckle patterns produced by a point source in a disordered medium. *Phys. Rev. A*, 82(4):043823, Oct 2010.
- [4] B. A. van Tiggelen and S. E. Skipetrov. Fluctuations of local density of states and c_0 speckle correlations are equal. *Phys. Rev. E*, 73:045601(R), apr 2006.
- [5] B. Shapiro. New Type of Intensity Correlation in Random Media. *Phys. Rev. Lett.*, 83:4733–4735, 1999.
- [6] R. Pierrat and R. Carminati. Spontaneous decay rate of a dipole emitter in a strongly scattering disordered environment. *Phys. Rev. A*, 81(6):063802, 2010.