The Legacy of E. Zavattini

Dimitar Bakalov

Institute for Nuclear Research and Nuclear Energy - Sofia - Bulgaria

This is an Open Access article distributed under the terms of the Creative Commons Attribution License 2.0, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited

Article published online by [EDP Sciences](http://www.edpsciences.org) and available at <http://www.iesc-proceedings.org> or <http://dx.doi.org/10.1051/iesc/2012qed01003>

Cargèse International School on QED, & Quantum Vacuum, Low Energy Frontier - 2012

2005: QED, Quantum Vacuum and the
Search for New Forces, Les Houches

2000: Frontier tests of QED and physics of the
vacuum, Trieste

1998: Frontier tests of QED and physics of the
vacuum, Sandansky

1994: Workshop on physics of exotic atoms, Erice

The Legacy of E.Zavattini

The Legacy of E.Zavattini

In the 1980's, in Dubna:

muon physics research, related to
“muon catalyzed fusion”.

E.Zavattini had already measured:

- muon capture rate
- muon life time
- cross sections with muonic atoms

The Legacy of E.Zavattini

- Precision experiments, which:
 - test fundamental theories
 - are aimed at solving key problems
 - display the interplay of weak, strong and EM interactions
 - consider alternative models

Precision experiments

Concluding remarks at the International Conference on a European Hadron Facility, Mainz 1986:

1. Muon capture rate and weak lepton-hadron form factors g_A and g_P
2. BNL proposal on the anomalous magnetic moment of the muon a_μ .

Testing QED

Particular interests:

Vacuum polarization and radiative effects:

- exotic atom spectroscopy
- muon anomaly measurements
- nonlinear QED effects

Of the Legacy of E.Zavattini:

Three selected topics:

- Nuclear muon capture
- Nonlinear effects of QED
- Exotic atom spectroscopy

Muon capture by nuclei

OMC: $\mu^- + p \rightarrow n + \nu_\mu$

$$\mu^- + d \rightarrow n + n + \nu_\mu$$

6 form factors at $q^2 \sim 0.88m_\mu^2$

$g_V(q^2)$, $g_M(q^2)$, $g_S(q^2)$, $g_A(q^2)$,
 $g_T(q^2)$, $g_P(q^2)$

Muon capture in H gas

First measurement of $\mu^- + p \rightarrow n + \nu_\mu$

PHYSICAL REVIEW

VOLUME 177, NUMBER 5

25 JANUARY 1969

Muon Capture in Gaseous Hydrogen

The experimental result is $\Lambda_{\text{expt}} = 651 \pm 57 \text{ sec}^{-1}$, which has to be compared with the theoretical rate $\Lambda_{s, \text{theor}} = 626 \pm 26 \text{ sec}^{-1}$. From the experimental capture rate, and within the framework of the currently accepted theory, we have obtained for the induced pseudoscalar coupling constant $g_p = (-7.3 \pm 3.7)g_V$. The results of the present experiment are analyzed, together with results obtained from stopping negative muons in liquid hydrogen.

To compare with: $\Lambda_S = 725.0 \pm 17.4$, $g_p = 7.3 \pm 1.1$
(MuCAP experiment, 2007)

Muon capture in liquid H

Muon capture in liquid H

Nuclear Physics A352 (1981) 365–378 © North-Holland Publishing Co., Amsterdam

A NOVEL MEASUREMENT OF THE MUON CAPTURE RATE IN LIQUID HYDROGEN BY THE LIFETIME TECHNIQUE

Abstract: The muon nuclear capture rate by protons has been measured by comparing the lifetime τ_{μ^-} of negative muons stopped in liquid hydrogen with the lifetime τ_{μ^+} of positive muons. We get $\tau_{\mu^-} = 2194.903 \pm 0.066$ ns and $\tau_{\mu^+} = 2197.182 \pm 0.121$ ns. Taking into account also the previous determinations of τ_{μ^+} , we present the updated determination of the muon lifetime.

Improved to $\tau_{\mu^+} = 2197.078(0.073)$ ns [PLB137(1984)]

To compare with MuLAN (2007), similar method used:

$$\tau_{\mu^+}(\text{MuLan}) = 2196980.3(2.2) \text{ ps.}$$

Ortho-para transitions in liquid H

Volume 104B, number 4

PHYSICS LETTERS

3 September 1981

MEASUREMENT OF THE ORTHO-PARA TRANSITION RATE IN THE $p\mu p$ MOLECULE AND DEDUCTION OF THE PSEUDOSCALAR COUPLING CONSTANT g_p^μ

The ortho-para transition rate in the $p\mu p$ molecule has been found experimentally to be $\lambda_{Op} = (4.1 \pm 1.4) \times 10^4 \text{ s}^{-1}$. Our recent result for the muon capture rate in liquid hydrogen can now be interpreted to extract the ortho-molecular capture rate: $\lambda_{OM} = (531 \pm 33) \text{ s}^{-1}$. A deduction of the pseudoscalar coupling constant g_p^μ is presented: we find $g_p^\mu = 8.7 \pm 1.9$.

Theoretical calculations: $\lambda_{op} = 7.1 \pm 4.1 \times 10^4 \text{ s}^{-1}$
Still an open problem.

Nonlinear QED effects

Vacuum birefringence in strong B:

$$n_{\perp} \neq n_{\parallel}$$

Leading to ellipticity and rotation of
linearly polarized light

(Analogous to Cotton-Mouton effect)

Vacuum birefringence in B

Euler-Heisenberg theory:

$$L = (E^2 - B^2)/(8\pi) + (A/4\pi)((E^2 - B^2)^2 + 7(E \cdot B)^2), \quad A = O(\alpha^2)$$

+h.o.t.

Experimental method

Volume 85B, number 1

PHYSICS LETTERS

30 July 1979

EXPERIMENTAL METHOD TO DETECT THE VACUUM BIREFRINGENCE INDUCED BY A MAGNETIC FIELD

E. IACOPINI and E. ZAVATTINI

CERN, Geneva, Switzerland

Received 28 May 1979

In this letter a method of measuring the birefringence induced in vacuum by a magnetic field is described: this effect is evaluated using the non-linear Euler–Heisenberg–Weisskopf lagrangian. The optical apparatus discussed here may detect an induced ellipticity on a laser beam down to 10^{-11} .

Vacuum birefringence in B

Other possible
sources: axions

From: Phys.Rev.D47(1993)3707 [left], Nucl.Instrum.Meth. A461(2001)329 [right]

Search for "axions" (BFRT)

PHYSICAL REVIEW D

VOLUME 47, NUMBER 9

1 MAY 1993

ARTICLES

Search for nearly massless, weakly coupled particles by optical techniques

Search for "axions" (PVLAS)

From: Phys.Rev.D77(2008)032006

Vacuum birefringence in B

- Vacuum birefringence in strong field: not yet observed
- Improved sensitivity experiments: in progress (BMV,...)
- Search for weakly interacting nearly massless particles: in progress
- Photon regeneration experiments...

Exotic atom spectroscopy

The electron replaced by a heavy charged particle.

$r \sim a_0 / (m/m_e)$: enhanced nucl.str.eff.

$E \sim E_0 * (m/m_e)$: different freq. range

- Kaonic atoms (DEAR)
- Antiprotonic atoms (Obelix)
- Muonic atoms (CERN, SIN)

Vacuum polarization in $(\mu\text{He})^+$

$2S_{1/2}-2P_{3/2}$ level difference

Nuclear Physics A278 (1977) 381 – 386;

PRECISE MEASUREMENT OF THE $2S_{1/2}-2P_{3/2}$ SPLITTING IN THE $(\mu^-{}^4\text{He})^+$ MUONIC ION

Abstract: The results of a new measurement of the $2S_{1/2}-2P_{3/2}$ splitting S^1 in the muonic ion $(\mu^-{}^4\text{He})^+$ are presented. We found $S_{\text{exp}}^1 = 1527.5 \pm 0.3 \text{ meV}$. Using the new, recently determined, value of the rms charge radius for ${}^4\text{He}$ we obtain for the difference D , between S_{exp}^1 and the corresponding theoretical prediction, the value $D = 0.2 \pm 4.2 \text{ meV}$: this value directly confirms, assuming μ -e universality, the QED vacuum polarization prediction to 0.25 %.

NOT reproduced by P.Hauser et al., PRA(1992) (low P)

Vacuum polarization in $(\mu\text{He})^+$

$2S_{1/2}-2P_{1/2}$ level difference

Volume 73B, number 2

PHYSICS LETTERS

MEASUREMENT OF THE $2S_{1/2}-2P_{1/2}$ SPLITTING IN THE $(\mu^{-4}\text{He})^+$ MUONIC ION

The measurement of the $2S_{1/2} \rightarrow 2P_{1/2}$ energy transition in muonic helium is presented. The energy difference S^1 is found to be $S_{\text{exp}}^1 = 1381.3 \pm 0.5 \text{ meV}$. This result agrees with the expected value $S^1 = 1381.2 \pm 0.3 \text{ meV}$ obtained assuming the previously measured value for the $2S_{1/2} \rightarrow 2P_{3/2}$ energy difference.

Confirmed by theory: 1381.716 meV (Martynenko 2006)

Metastability of $(\mu\text{He})^+_{2S}$ at 40 A?

VOLUME 33

29 JULY 1974

NUMBER 5

Measurement of the Initial Population and Decay Rate of the $(\mu^4\text{He})_{2S}^+$ System
in a Helium Target at 50 atm

Experimentally observed:

2S-fraction: $\sim 4\%$

2S-lifetime: $\sim 1\mu\text{s}$ at 50 At.

Metastability of $(\mu\text{He})^+_{2s}$ at 40 A?

PHYSICAL REVIEW A

VOLUME 41, NUMBER 5

**Formation and disappearance rates of metastable muon- α levels
in high-pressure helium targets**

Semi-quantitative theoretical model:
prompt formation of excited clusters
 $[(\mu^-\text{He})_{2s}\text{-He}]^*$, $[(\mu^-\text{He})_{2s}\text{-He-He}]^*$, ...

Metastability of $(\mu\text{He})^+_{2s}$ at 40 A?

But:

M.Eckhause et al., PRA33(1986) found:
at 40 At, the lifetime of $(\mu\text{He})^+_{2s}$ is 40 ns.

2S-metastability: an open problem.

Future experiments expected (PSI?)

Born-Infeld theory (1983)

IL NUOVO CIMENTO

VOL. 78 B, N. 1

11 Novembre 1983

Vacuum Polarization Effects in the $(\mu^{-4}\text{He})^{+}$ Atom and the Born-Infeld Electromagnetic Theory.

$$L = b^2 \left(1 - \sqrt{1 - (E^2 - B^2)/b^2 - (E \cdot B)^2/b^4} \right)$$

Lamb shift in $(\mu\text{He})^{+}$ can be reproduced by adjusting b

No vacuum birefringence produced

Antiprotonic helium spectroscopy (Obelix Collaboration)

- Delayed annihilation of antiprotons stopped in Helium
- Dependence on contaminants
- Alternative mechanism suggested:
formation of $[(p\text{-He})^+\text{He}]^*$
(not confirmed)

Muonic hydrogen spectroscopy

- Elastic scattering cross sections
- Muon transfer to heavier gases (xenon)
- Laser spectroscopy in excited states
- Heavier elements muonic atoms

Hyperfine splitting in $(\mu\text{H})_{1s}$?

HFS in hydrogen-like atoms

$$\Delta E = \Delta E^F (1 + \delta^{\text{QED}} + \delta^{\text{str}})$$

$$\delta^{\text{str}} = \delta^{\text{rigid}} + \delta^{\text{pol}} + \delta^{\text{hvp}} + \dots$$

$$\delta^{\text{rigid}} = \delta^{\text{Zemach}} + \delta^{\text{recoil}}$$

$$\delta^{\text{Zemach}} = -2\alpha m R_p + O(\alpha^2)$$

$$R_p = \langle r \rangle_{E^0 M}; R_p^2 \neq r_{\text{sq}}^2$$

$$\Delta^{\text{pol}} \leftrightarrow \text{polarizability 'a' ?}$$

Two parameters (a , R_p) from measurements in H & μH ?

Hyperfine splitting in $(\mu\text{H})_{1\text{S}}$

- Time distribution of the events of μ^- transfer to:
 - walls of target volume
 - nuclei of appropriate heavier gas admixture
- Tunable IR laser source

Nuclear Instruments and Methods in Physics Research A 341 (1994)
Compact waveguide FEL for spectroscopic measurements in muonic hydrogen

From: Hyperf.interact.136(2001)1

The Legacy of E. Zavattini

Ongoing experimental activities

- MuLAN & MuCAP
- Vacuum birefringence
- Muonic helium spectroscopy
- Muonic hydrogen hyperfine structure

...